



IJABBR- 2014- eISSN: 2322-4827

International Journal of Advanced Biological and Biomedical Research

Journal homepage: [www.ijabbr.com](http://www.ijabbr.com)



## Original Article

### Effect of Acyclovir on the Viability of Chicken Embryo During the Developmental Stage

Seyede Saeedeh Mosallanejad<sup>1\*</sup>, Sajedeh Salandari<sup>2</sup>, Hadi Tavakkoli<sup>3</sup>

<sup>1</sup>Graduate student, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

<sup>2</sup>Graduate student, Faculty of Veterinary Medicine, Shahid Bahonar University of Kerman, Kerman, Iran

<sup>3</sup>Department of Clinical Science, Faculty of Veterinary Medicine, Shahid Bahonar University of Kerman, Kerman, Iran

#### ARTICLE INFO

##### Article history:

Received: 08 Feb, 2015

Revised: 15 Mar, 2015

Accepted: 25 Apr, 2015

ePublished: 30 May, 2015

##### Key words:

Acyclovir

Chicken

Embryo

Pathology

#### ABSTRACT

**Objective:** In recent years, antiviral drugs have been used widely in the globe to prevent and treat viral diseases. They injected into eggs to eliminate pathogens and prevent of egg transmission of disease, but the adverse effects of drugs have always been a major concern. There is little information available about the safety of antiviral drugs in the embryonated eggs of birds. The objective of this study was to evaluation of the effect of acyclovir on the viability of chicken embryo during the first trimester of the incubation period. **Methods:** Fertile chicken eggs were divided into two equal treatment groups. The phosphate buffered saline injected group and acyclovir-injected group whose individuals were injected with acyclovir solution at a dosage of 80 mg per Kg egg-weight. Embryos monitored daily during incubation until day 10 after which; they were examined for viability. **Results:** Results showed that all of the embryos were dead in the acyclovir-injected group. Based on findings, it is concluded that acyclovir at above-mentioned concentration has a severe lethal effect on the chicken embryo. So, acyclovir egg-injection cannot be used to eliminate pathogens and prevention of egg transmission of the disease.

#### 1.INTRODUCTION

Raising chickens for their eggs and meat is becoming a popular and lucrative business in many countries including Iran. Viral diseases are the most common and destructive disease of the chicken (Swayne et al. 2013). So, different types of antiviral drugs were used to prevent and treatment of viral diseases in this industry (Swayne, Glisson et al. 2013; Tavakkoli et al. 2104).

Acyclovir belongs to the guanosine analogue antiviral medication and is on the World Health Organization's List of Essential Medicines. It is primarily used for the treatment of herpes simplex virus infections, chickenpox and shingles (Tavakkoli et al. 2014a). Other uses

including prevention of cytomegalovirus infections following transplant, infections due to Epstein-Barr virus, genital herpes simplex, cold sores, shingles, acute chickenpox in immunocompromised patients, herpes simplex encephalitis, acute mucocutaneous HSV infections in immunocompromised patients and herpes of the eye and herpes simplex blepharitis (Richardson 1997; Sweetman et al. 2009; Tavakkoli et al. 2014c; Tavakkoli, Derakhshanfar et al. 2104). Acyclovir trials show that this agent has no role in preventing HIV transmission, but it can help slow HIV disease progression in people not taking anti-retroviral therapy (Richardson 1997; Conway et al. 2013; Wightwick et al. 2013). In veterinary medicine, acyclovir is used to prevent disease and cure animals and birds (Ahrens et al.

\*Corresponding Author: Seyede Saeedeh Mosallanejad, Graduate student, School of Veterinary Medicine, Shiraz University, Shiraz, Iran ([Mosala.sm@hotmail.com](mailto:Mosala.sm@hotmail.com))

2013; Swayne, Glisson *et al.* 2013). In hatcheries, the hygienic process in association with injecting drugs into the egg, result in eliminating infection and preventing egg transmission of pathogens. Alternatively, eggs may be dipped in the drug solutions for controlling the disease transmission before hatching.

Adverse effects of drugs have always been a major concern. There is little research in the literature describing the effect of acyclovir on the viability of bird embryos, and further studies still need to be undertaken to determine this aspect. In this regard, in the present study, we evaluated the effect of acyclovir on the viability of the chicken embryo during the first trimester of the incubation period. We believe that results in this study will contribute to our better understanding of the safety of antiviral drugs for *in ovo* administration in the bird's egg.

## 2. MATERIALS AND METHODS

### 2.1. Drug and eggs

Acyclovir powder was diluted in phosphate buffered saline solution. A volume of 0.5 mL of phosphate buffered saline solution with 80 mg acyclovir was inoculated per Kg egg-weight. Fertile chicken eggs (Ross 308) with the average egg-weight of  $50 \pm 1$  g and with the same age were purchased from a local breeder farm. In this farm, birds were kept and grown up under the standard condition of breeding.

### 2.2. Experimental protocol


Eggs were incubated at 37.5°C and 55% relative humidity. The eggs were randomly assigned to two equal treatment groups, 10 eggs each, as follows. Group 1: phosphate buffered saline injected group, embryonated eggs injected with sterile phosphate buffered saline of 0.5 ml/egg into the yolk sac. On day 4 of incubation, the eggs of group 2 treated with acyclovir solution at a dosage of 80 mg acyclovir per Kg egg-weight. Embryos received treatment by direct injection into the yolk sac according to the standard techniques (Hamburger 1942). Embryos were re-incubated post-treatment and allowed to develop. The viability of the embryos was checked throughout the incubation period by candling. At the end of the experiment, on day 10, the eggs were opened at the wider end and monitored for the viability of the embryos. The treatment protocols and procedures in this study were conducted according to local ethical guidelines, and were approved by the Animal Ethics Committee of the Research Council of Shahid Bahonar University, Iran.

### 2.3. Statistical analysis


Statistical analysis was performed using SPSS version 20. The Chi-Square test was used to determine the significant differences in embryo-viability between experimental groups. A P-value of  $<0.05$  was considered as statistically significant.

## 2. RESULTS

Candling of the hatching chicken eggs during the incubation period and examination of the eggs or embryos is a useful tool for hatchery managers. In the current study, the developing blood vessels were seen in the embryonated eggs at the early stage of incubation (figure 1). In the next phase of incubation, the more advanced circulatory system was not developed. Results of our experiments showed that the early developing blood vessels were seen in 80% and 70% of chicken eggs received phosphate buffered saline and treated with acyclovir, respectively. At the end of the experiment, on day 10, the eggs were opened at the wider end and monitored for the viability of the embryos. As can be seen in the figure 2, there was centrally located dead embryo in the chicken egg treated with acyclovir into the yolk sac. The circulatory system and the blood vessels of the extra-embryonic membranes were not developed, while in the chicken egg treated with phosphate buffered saline, the embryo and blood vessels were normal.



**Fig. 1.** The developmental stage of a normal chicken embryo. The embryo and developing blood vessels are seen (arrows).



**Fig. 2.** The chicken egg treated with acyclovir into the yolk sac. A centrally located dead embryo is seen.

## 4. DISCUSSION

Despite reports of an expanding chicken industry across the globe, pathogenic agents can decrease the hatchability rate during embryonic development.

Embryonic death causes serious economic losses to the poultry industry (Swayne, Glisson *et al.* 2013).

For many years, researchers have been using different antipathogenic compounds to restrict pathogens and enhance the performance of different poultry species, including young chicken (Colomer-Lluch *et al.* 2011; Sapkota *et al.* 2011; Obeng *et al.* 2012; Banerjee *et al.* 2013), chicken (McDougald *et al.* 2012; Crespo *et al.* 2013; Rigobelo *et al.* 2013), turkey (Altunsoy *et al.* 2011; Erdem *et al.* 2012; Buscaglia 2013), broiler (MacDonald *et al.* 2011; Agunos *et al.* 2012; Lee *et al.* 2012; Tavakkoli *et al.* 2014b; Tavakkoli, Derakhshanfar *et al.* 2014c), layers (Hasan *et al.* 2011; Lee *et al.* 2013; Nemati 2013) and poultry breeder (Kabir 2010; Priyantha *et al.* 2012; Jones *et al.* 2013; Tavakkoli, Derakhshanfar *et al.* 2104).

Antiviral drugs have an increased role as therapeutic agents against avian viruses. They have virucidal effect and a wide antiviral spectrum. Most viruses are susceptible (Sweetman, Pharm *et al.* 2009; Ahrens and Martin 2013). Acyclovir belongs to the guanosine analogue antiviral group. It has been used successfully for several decades in many countries such as Canada, Spain, France, Austria, Polish, Denmark, Germany, Turkey, Africa, United States and China. In recent years, its use has increased rapidly in the Iranian pet bird industry, but there is little information available about the effects of injecting the acyclovir solution into the bird's egg. In the present study, we have evaluated the effect of acyclovir on the viability of the chicken embryo during the first trimester of the incubation period.

Up to now, drug-egg-treatment has been examined and described in different situations (Ghazikhanian *et al.* 1980; Sheeks *et al.* 1992; Kleven 2008; Singroha *et al.* 2012; Singroha *et al.* 2013; Tavakkoli *et al.* 2013; Tavakkoli, Derakhshanfar *et al.* 2014b). The results of these studies show that injecting drugs into hatching eggs can eliminate pathogens and prevent vertical transmission of disease. Some injection sites that are present in fertile eggs at day 4 of incubation are the air cell and yolk sac. Our results obviously showed severe lethal effect was seen in the chicken eggs treated with acyclovir solution by the yolk sac rout. Therefore, these findings suggest that acyclovir is not safe for injection into the ovo by the yolk sac route. Nevertheless, further efforts are needed to evaluate in ovo administration of various antiviral drugs for prevention and eliminate pathogenic microorganisms. In conclusion, based on findings, it is concluded that acyclovir cannot be used for the success of the prevention and treatment scheme in chicken embryo.

### Acknowledgment

Authors wish to thank Dr. A. Hamidi for improving the use of English in the manuscript.

### REFERENCES

- Agunos A, Léger D, Carson C (2012). Review of antimicrobial therapy of selected bacterial diseases in broiler chickens in Canada. *Canadian Vet. J.* 53(12): 1289.
- Ahrens FA, Martin RJ (2013). Antimicrobial drugs. *Handbook of Veterinary Pharmacology* 347.
- Altunsoy A, Aypak C, Azap A, Ergönül Ö, Balık İ (2011). The impact of a nationwide antibiotic restriction program on antibiotic usage and resistance against nosocomial pathogens in Turkey. *Int. J. Med. Sci.* 8(4): 339.
- Banerjee S, Mukhopadhyay SK, Ganguly S (2013). Phytogetic Growth Promoter as Replacers for Antibiotic Growth Promoter in Poultry Birds. *J. Anim. Genet. Res.* 1(1): 6-7.
- Buscaglia C (2013). Influence of the addition of antibiotics on survival of herpesvirus of turkeys. *Avian Dis.* 52(2): 437-436.
- Colomer-Lluch M, Imamovic L, Jofre J, Muniesa M (2011). Bacteriophages carrying antibiotic resistance genes in fecal waste from cattle, pigs, and poultry. *Antimicrob. Agents Chemother.* 55(10): 4908-4911.
- Conway GR, Pretty JN (2013). *Unwelcome harvest: agriculture and pollution*, Routledge.
- Crespo R, Shivaprasad H, Silva Franca M (2013). Ulcerative Enteritis-like Disease Associated with *Clostridium sordellii* in Quail. *Avian Dis.* 57(3): 213-221.
- Erdem H, Akova M (2012). Leading infectious diseases problems in Turkey. *Clin. Microbiol. Infect.* 18(11): 1056-1067.
- Ghazikhanian GY, Yamamoto R, McCapes R, Dungan WM, Larsen C, Ortmayer H (1980). Antibiotic Egg Injection to Eliminate Disease II. Elimination of *Mycoplasma meleagridis* from a Strain of Turkeys. *Avian Dis.* 48-56.
- Hamburger V (1942). *A manual of experimental embryology*, University of Chicago Press Chicago.
- Hasan B, Faruque R, Drobni M, Waldenström J, Sadique A, Ahmed KU, Islam Z, Parvez MH, Olsen B, Alam M (2011). High prevalence of antibiotic resistance in pathogenic *Escherichia coli* from large-and small-scale poultry farms in Bangladesh. *Avian Dis.* 55(4): 689-692.
- Jones K, Thornton J, Zhang Y, Mauel M (2013). A 5-year retrospective report of *Gallibacterium anatis* and *Pasteurella multocida* isolates from chickens in Mississippi. *Poult. Sci.* 92(12): 3166-3171.

Kabir S (2010). Avian colibacillosis and salmonellosis: a closer look at epidemiology, pathogenesis, diagnosis, control and public health concerns. *Int. J. Environ. Res. Public Health* 7(1): 89-114.

Kleven S (2008). Control of avian mycoplasma infections in commercial poultry. *Avian Dis.* 52(3): 367-374.

Lee K-W, Ho Hong Y, Lee S-H, Jang SI, Park M-S, Bautista DA, Donald Ritter G, Jeong W, Jeoung H-Y, An D-J (2012). Effects of anticoccidial and antibiotic growth promoter programs on broiler performance and immune status. *Res. Vet. Sci.* 93(2): 721-728.

Lee S-K, Chon J-W, Song K-Y, Hyeon J-Y, Moon J-S, Seo K-H (2013). Prevalence, characterization, and antimicrobial susceptibility of *Salmonella Gallinarum* isolated from eggs produced in conventional or organic farms in South Korea. *Poult. Sci.* 92(10): 2789-2797.

MacDonald JM, Wang S-L (2011). Foregoing sub-therapeutic antibiotics: The impact on broiler grow-out operations. *App. Economic Perspectives Policy* 33(1): 79-98.

McDougald L, Abraham M, Beckstead R (2012). An Outbreak of Blackhead Disease (*Histomonas meleagridis*) in Farm-Reared Bobwhite Quail (*Colinus virginianus*). *Avian Dis.* 56(4): 754-756.

Nemati M (2013). Antimicrobial resistance of *Porteus* isolates from poultry. *Europ. J. Experiment. Biol.* 3(6): 499-500.

Obeng AS, Rickard H, Ndi O, Sexton M, Barton M (2012). Antibiotic resistance, phylogenetic grouping and virulence potential of *Escherichia coli* isolated from the faeces of intensively farmed and free range poultry. *Vet. Microbiol.* 154(3): 305-315.

Priyantha M, Vipulasiri A, Gunawardana G (2012). *Salmonella* control in poultry breeder farms in Sri Lanka: Effects of oral antibiotic treatment on whole blood agglutination test with *Salmonella pullorum* antigen. *Int. J. Livestock Product.* 3(2): 21-24.

Richardson HW (1997). *Handbook of copper compounds and applications*, CRC Press.

Rigobelo EC, Blackall PJ, Maluta RP, Ávila FAd (2013). Identification and antimicrobial susceptibility patterns of *Pasteurella multocida* isolated from chickens and Japanese quails in Brazil. *Braz. J. Microbiol.* 44(1): 161-164.

Sapkota AR, Hulet RM, Zhang G, McDermott P, Kinney EL, Schwab KJ, Joseph SW (2011). Lower prevalence of antibiotic-resistant enterococci on US conventional poultry farms that transitioned to organic practices. *Environ. Health Perspect.* 119(11): 1622.

Sheeks OB, Sheeks RL (1992). Egg injection method, apparatus and carrier solution for improving hatchability and disease control, Google Patents.

Singroha R, Srivastava S, Chhikara P (2012). Effect of Gentamicin on kidney in developing chicks. *Eur J Anat* 16(2): 119-126.

Singroha R, Srivastava S, Chhikara P (2013). Effect of gentamicin on proximal convoluted tubules of kidney in developing chicks. *J. Anat. Soc. India* 62(1): 17-22.

Swayne D, Glisson JR, McDougald L, Nolan LK, Suarez DL, Nair VL (2013). *Diseases of Poultry*, Wiley-Blackwell.

Sweetman SC, Pharm B, PharmS F, Eds. (2009). *Martindale: The Complete Drug Reference*. London, Pharmaceutical Press.

Tavakkoli H, Derakhshanfar A, Noori Gooshki S (2013). A short preliminary experimental study on teratogenic effect of methenamine in embryonic model. *International Journal of Advanced Biological and Biomedical Research.* 1(12): 1523-1528.

Tavakkoli H, Derakhshanfar A, Noori Gooshki S (2014a). The effect of florfenicol egg-injection on embryonated chicken egg. *International Journal of Advanced Biological and Biomedical Research.* 2(2): 496-503.

Tavakkoli H, Derakhshanfar A, Noori Gooshki S (2014b). Toxicopathological lesions of fosfomycin in embryonic model. *Euro. J. Experiment. Biol.* 4(2): 63-71.

Tavakkoli H, Derakhshanfar A, S S (2014c). Toxicology of urotropine in chicken embryo model. *Online J Vet Res* 18(2): 109-115.

Tavakkoli H, Derakhshanfar A, Salandari S (2104). Investigation on the using of linco-spectin solution for in ovo administration in chicken embryo. *International Journal of Advanced Biological and Biomedical Research.* 2(1): 110-116.

Wightwick AM, Salzman SA, Reichman SM, Allinson G, Menzies NW (2013). Effects of copper fungicide residues on the microbial function of vineyard soils. *Environment. Sci. Pollution Res.* 20(3): 1574-1585.